



Bacteriological Assessment of Smoked-Dried White Shrimp (*Nematopalaemon hastatus* Aurivillius, 1898) Sold in Calabar, Nigeria

Tarh, Jacqueline Ebob^a, Bassey, Emmanuel Effiong^{b#*}, Mbah, Maurice^b, Ekpiken, Ekpiken Solomon^a and Upula, Sylvanus Akpak^a

^a Department of Microbiology, Faculty of Biological Sciences, Cross River University of Technology, Calabar, Nigeria.

^b Department of Medical Laboratory Science, Faculty of Allied Medical Sciences, College of Medical Sciences, University of Calabar, Calabar, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2023/v15i41304

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97614>

Original Research Article

Received: 22/01/2023

Accepted: 24/03/2023

Published: 13/04/2023

ABSTRACT

Aim: Enumeration of bacterial counts is an important index of assessing the safety and quality of food products. This study aimed to investigate the occurrence of pathogenic bacteria in white shrimp samples obtained from major markets serving consumers in Calabar.

<https://orcid.org/0000-0003-2029-5611>

*Corresponding author: Email: emmanueleffiong2046@gmail.com;

Study Design: This study was a cross sectional study conducted between April 2017 and April 2018.

Methodology: One hundred and twenty (120) samples of smoked dried *Nematopalaemon hastatus* (white shrimp) were analyzed for the presence of bacterial pathogens. Bacterial loads and identification of isolated organisms was determined using standard microbiological methods.

Results: The results showed that 66.7% of the analyzed shrimps had aerobic bacterial counts exceeding the upper permissible limit ($<1.0 \times 10^6$ Cfu/g) and 56.7% had unsatisfactory (>20 Cfu/g) *Vibrio* counts. The study revealed the presence of different bacteria genera namely *Klebsiella*, *Salmonella*, *Pseudomonas*, *Serratia*, *Vibrio*, *Citrobacter*, *Proteus*, *Aeromonas*, *Streptococcus*, *Escherichia*, *Coagulase-negative Staphylococci* (CoNS), *Enterobacter* and *Bacillus*. Predominant organism was *Salmonella* spp (26.7%), followed by *Vibrio* spp (21.7%) while the least isolated organisms were *Bacillus* spp and *Coagulase-Negative Staphylococci* (1.7%) each. The occurrence of high counts of pathogens in seafood may cause food poisoning; especially in individuals who consume this seafood raw, or lightly or insufficiently cooked.

Conclusion: Hence, seafood should be processed and packaged under standard hygienic conditions to reduce the risk of microbial contamination. In addition, public health awareness campaign targeted at consumers and vendors should be optimized with frequent monitoring by regulatory agencies.

Keywords: Bacteriological assessment; smoked white shrimps; Calabar; Nigeria.

1. INTRODUCTION

White shrimp (*Nematopalaemon hastatus*, Aurivillius, 1898) is one of the most important commercially valuable seafood consumed not only in Nigeria but also in sub-Saharan Africa and other regions of the world. Due to its nutritional benefits, they are consumed daily in soups and other recipe among coastal and inland dwellers [1].

Seafood is an important source of animal protein, omega-3 fatty acids, essential amino acids, essential micronutrients, vitamin and minerals [2,3]. Nutritionally, consumption of seafood is associated with potential health benefits including neurologic development during gestation and infancy, and reduced risk of heart disease [4]. Carbohydrate content in seafood is known to be very little and as such recommended for weight loss [5,3]. In Nigeria, white shrimps are often sold and consumed in dried form, however, most coastal dwellers consume it fresh [1].

Seafood constitutes over 40% of the animal protein consumed in Nigeria compared to meat and it is relatively less expensive. This accounts for the mass preference for fish products [6]. Despite the nutrients and benefits derived from seafood, consumption of contaminated seafood is accompanied with potential health risks [7]. Seafood are prone to microbial contamination, especially filter feeders such as shrimps, which concentrate these pathogens in their filtration

systems [7,8]. Infectious agents involved in seafood contamination include bacteria, fungi, viruses and parasites, which can cause illnesses ranging from mild gastroenteritis to life-threatening diseases [9]. Additionally, some of these pathogens are naturally present in the aquatic environment, while others can be introduced through animal or human fecal shedding and sewage pollution [4]. Additionally, seafood may become contaminated during handling, processing, or preparation. Furthermore, microbiological evaluation of shellfish has shown that they harbour many pathogenic microorganisms that have been implicated in outbreak of foodborne diseases in many part of the world including Nigeria, where it is highly consumed [10]. These illnesses include Typhoid fever, Cholera, diarrhoea and other digestive disorders [6,11].

Also, seafood provides a good medium for the proliferation of known medically important pathogens including *Escherichia coli*, [12] *Salmonella* species [2] *Vibrio* species, [7] *Pseudomonas* species etc. [13,6]. Although adequate cooking has been shown to inactivate most pathogenic contaminants in meat and poultry products, seafood is sometime consumed raw or with minimal cooking especially among coastal dwellers. Findings from previous studies have reported bacterial contaminants in white shrimps [9,14]. However, there are paucity of related studies investigating the contamination rate of white shrimps in Calabar. Consequently, this study seeks to analyze the occurrence of

pathogenic bacteria in white shrimp (*N. hastatus*) samples obtained from major markets serving consumers in Calabar.

2. MATERIALS AND METHODS

2.1 Study Area

This study was carried out in Calabar, a city in Cross River State, in Southern Nigeria. The city lies between Longitudes 4°57'0"N and 8°19'30"E and Latitudes 4°95'0N and 8°32'0"E. Calabar is divided administratively into Calabar Municipal and Calabar South LGAs. The city is mostly populated with Efik speaking people. The city has an area of 406 square kilometers and had a population of 371, 022 at the 2006 census. The study was conducted in two major markets (Watt market and Marian market) and two beaches (Esuk-Nsidung beach and Obufa Esuk beach). These sites were selected because they are the most visited market in Calabar metropolis [15].

2.2 Sampling Method

This study was a cross sectional study conducted between April 2017 and April 2018. A total of 120 white shrimps (Fig. 1) samples were purchased from 2 major market (i.e. Marian market and Watt market) and 2 beaches markets (i.e. Esuk-Nsidung beach market and Obufa Esuk beach market) in Calabar metropolis. Each sample was collected into a sterile polythene bag and sealed to avoid further contamination. All samples obtained were labeled appropriately by indicating the location, time and date of collection. The samples were subsequently conveyed for further evaluation and microbiological analysis. Ten (10 g) of each shrimp sample was homogenized using sterile laboratory mortar. About 1.0 g of the homogenate was mixed with 10 mL of alkaline peptone water and incubated at 37°C for about 6-7 hours. The suspension was serially diluted 10-fold using normal saline, and 0.5 mL of the appropriate dilution was spread over freshly prepared Thiosulfate Citrate Bile-Salt Sucrose (TCBS) Agar plate and MacConkey Agar plate, using a sterile spreader. The plates were incubated at 37°C for 18-24 hours. Following incubation, the plates were observed for characteristic colonies. Discrete representatives of each unique colony type were selected for purification, characterization and identification using conventional microbiological and biochemical tests.



Fig. 1. White shrimp (*N. hastatus*)

(Source: Emmanuel E. Bassey, 2023, Watt market)

2.3 Data Analysis

Data was analyzed using Statistical Package for the Social Sciences (SPSS) software version 20 and results presented as percentage. Samples contamination rate was interpreted using the guideline provided by Center of Food Safety [16].

3. RESULTS AND DISCUSSION

Enumeration of bacterial count is an important index of assessing the safety and quality of food products. Results obtained revealed that all analyzed *N. hastatus* samples were contaminated with various degree of bacterial growth (Table 1). According to the Center of Food Safety guideline [16], acceptable upper limit of aerobic bacterial count in shellfish is $<1.0 \times 10^6$ Cfu/g. The present study showed that 33.3% (40/120) of the examined shrimps had bacterial count under the permissive limit, 56.7% (68/120) were at borderline ($10^6 \leq 10^7$ Cfu/g) while 10.0% (12/120) of samples were unsatisfactory ($>10^7$ Cfu/g) for consumption. Of the 12 samples that were unsatisfactory, 8 (26.7%) were obtained from Nsidung beach market while 4 (13.3%) were from Obufa Esuk beach market. Previous studies have shown that high microbial loads can be obtained from different species of shrimps. Thus, Ajibare *et al.* [9] reported total heterotrophic bacterial counts (THBC) of 1.076 to 1.125×10^2 Cfu/g from white shrimp (*N. hastatus*) in Ondo state, Nigeria. Also, the high bacterial load observed in this study corroborated with Orji *et al.* [14] in Ebonyi state, Nigeria, who reported mean aerobic counts of $1.87 \pm 0.543 \times$

10^7 Cfug from fresh shrimps (*Palaemon serratus*). Similarly, another study conducted by Amin et al. [17] in Egypt recorded mean aerobic counts of $4.62 \times 10^4 \pm 1.7 \times 10^3$ Cfug. The results of the current study also agreed with Talukder et al. [18] in Bangladesh who recorded total viable counts between $\log 2.92 \pm 0.22$ Cfug and $\log 3.41 \pm 0.23$ Cfug from shrimps (*Penaeus monodon*) in Rupsha and Mongla respectively.

In this study, 56.7% (68/120) of the analyzed samples had *Vibrio* counts higher than the upper acceptable limit [16]. This aligns with finding by Talukder et al. [18] who reported *Vibrio* count ranging from $\log 2.06$ to 2.11 Cfug in shrimps (*P. monodon*) and Nwosu et al. [19] who reported heavy *Vibrio* counts in smoked dried shrimps (*Litopenaeus vannamei*) sold in markets compared to home-smoked dried rinsed shrimps.

The elevated aerobic bacterial counts recorded in this study could be attributed to factors such as the use of polluted water at the shores, poor handling, unhygienic environment where these fish products are sold, inappropriate storage and exposure to insect vectors [20,21,19]. Inadequate basic amenities such as potable water in major fishing communities in Nigeria remain a major challenge [22,19]. This has encouraged the use of polluted sea water in washing shrimps which inadvertently

predisposes shrimps to further microbial contamination and may have contributed significantly to the observed contamination rate.

From this study, thirteen different bacterial genera were recovered from *N. hastatus* (Fig. 2). The bacteria isolated include species of *Klebsiella*, *Salmonella*, *Pseudomonas*, *Serratia*, *Vibrio*, *Citrobacter*, *Proteus*, *Aeromonas*, *Streptococcus*, *Escherichia*, *Coagulase-negative Staphylococci* (CoNS), *Enterobacter* and *Bacillus*. The occurrence of these bacterial species is of public health concern because majority are known pathogen implicated in gastrointestinal disorder such as diarrhea, dysentery, typhoid fever, cholera among others [9]. These organisms have been previously reported as major contaminants of seafood [6,9,14]. This also is in agreement with Hariharan and Amadi [13] who reported the prevalence of pathogenic bacteria in shellfish.

This study indicates high occurrence of *Salmonella* spp (26.7%) and *Vibrio* spp (21.7%) in the analyzed samples which is of public health importance. Similar observation on the prevalence of *Salmonella* spp was reported by Bakr et al. [7] in Alexandria, Egypt. The potential source of *Salmonella* contamination in *N. hastatus* is likely due to poor water quality, surface water run-off and fecal contamination from wild animal or livestock [23].




Fig 2. Prevalence rate of the isolated bacteria

Table 1. Contamination rate of dried crayfish obtained in Calabar

Location	No of samples	Aerobic Bacterial Counts			Vibrio Counts		
		Satisfactory $1.0 \times 10^6 \text{Cfu/g}$	Borderline $10^5 - \leq 10^7 \text{Cfu/g}$	Unsatisfactory >math>10^7 \text{Cfu/g}</math>	Satisfactory 20Cfu/g	Borderline $20 - \leq 10^3 \text{Cfu/g}$	Unsatisfactory >math>10^3 \text{Cfu/g}</math>
Watt market	30	10(33.3)	20(66.7)	0(0.0)	16(53.3)	8(26.7)	6(20.0)
Marian market	30	20(66.7)	10(33.3)	0(0.0)	12(40.0)	6(20.0)	12(40.0)
Nsi-dung beach	30	8(26.7)	14(46.7)	8(26.7)	6(20.0)	20(66.7)	4(13.3)
Obufa Esuk beach	30	2(6.7)	24(80.0)	4(13.3)	18(60.0)	10(33.3)	2(6.7)
Total	120	40(33.3)	68(56.7)	12(10.0)	52(43.3)	44(36.7)	24(20.0)

The occurrence of *Escherichia coli* in seafood is considered a sanitary case and may represent a health risk to consumers if related to pathogenic strains, especially diarrheagenic *Escherichia coli*. However, the presence of non-pathogenic *Escherichia coli* in seafood is a concern to public health, since this bacterium is recognized as an indicator of fecal contamination, possibly indicating the presence of other enteric pathogens [12].

The presence of *Pseudomonas* species in the analyzed samples is worrisome. *Pseudomonas* species are implicated in Pseudomonal food poisoning. These infections are complicated and life threatening [24]. *Citrobacter*, *Proteus*, *Serratia*, *Klebsiella* and *Enterobacter* are mostly members of the *Enterobacteriaceae* family which are mainly fecal contaminants that could be picked from unhygienic sources and they are implicated in gastrointestinal disorder etc. [9]. The disease caused by *Staphylococcus aureus* is foodborne intoxication and common symptoms include nausea, but in severe cases dehydration can lead to shock and collapse [6]. *Aeromonas* spp is ubiquitous in fresh water environment, and their presence is in consonance with Markey et al. [25]. The presence of CoNS implies contamination from human during handling [14, 18].

The survival of any microbial group, within a particular ecological niche, is greatly dependent on environmental parameters and anthropogenic activities [26]. Hence, abundance of plethora of microbial groups in the aquatic environment have a significant effect in the high contaminated rate [22]. The differences in the prevalence of isolated organisms between this and earlier studies could be explained when factors such as differences in sample size, storage condition, hygiene, environmental and geographical variation are considered.

4. CONCLUSION

White shrimps, like any other seafood has the potential of causing foodborne illnesses from bacterial, fungal, viral and parasitic pathogens under certain conditions. Hence, it is encouraged that the conditions of processing and packaging of seafood be done under standard hygienic measures to reduce the risk of microbial contamination. In addition, public health awareness campaign targeted at consumers and vendors should be optimized with frequent monitoring by regulatory agencies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bayode PO, Olabode TA, Taiwo TA. Consumers' preference and perception of smoke-dried white shrimp (*Nematopalaemon hastatus*) in coastal areas of Ondo State, Nigeria. Int J Fish Aquacult. 2018;10(3):22-33. DOI: 10.5897/IJFA2017.0661.
2. Amagliani G, Brandi G, Schiavano GF. Incidence and role of Salmonella in sea food safely. Food Res Int. 2012;45(2):780-8.
3. Venugopal V, Gopakumar K. Shellfish: nutritive value, health benefits, and consumer safety. Compr Rev Food Sci Food Saf. 2017;16(6):1219-42. DOI: 10.1111/1541-4337.12312
4. Iwamoto M, Ayers T, Mahon BE, Swerdlow DL. Epidemiology of seafood-associated infections in the United States. Clin Microbiol Rev. 2010;23(2):399-411. DOI: 10.1128/CMR.00059-09
5. Elegbede IO, Fashina-Bombata HA. Proximate and mineral compositions of common crab species [*Callinectes pallidus* and *Cardisoma armatum*] of Badagry Creek, Nigeria. Poutl Fish Wildl Sci. 2013;2:110. DOI: 10.4172/pfw.1000110.
6. Nrior RR, Iyibo SN, Ngerebara NN. Microbiological assessment of niger delta shell sea foods; periwinkle (*Tympanotonus Fuscatus*), Oyster (*Crassostrea virginica*) and Veined rapa whelk (*Rapana venosa*) from crude oil polluted site. Int J Curr Res Multidiscip (IJCRM). 2016;2(7):01-9.
7. Bakr WMK, Hazzah WA, Abaza AF. Detection of Salmonella and Vibrio spp in some sea food in Alexandria. J Am Sci. 2011;7(9):663-7.
8. Topic Popovic N, Benussi Skukan A, Dzidara P, Coz-Rakovac R, Strunjak-Perovic I, Kozacinski L et al. Microbiological quality of marketed fresh and frozen seafood caught off the Adriatic coast of Croatia. Vet Med. 2010;55(5):233-41.
9. Ajibare AO, Olawusi-Peters OO, Oyinlola KA. Microbiological load evaluation of white shrimp (*Nematopalaemon hastatus* Aurivillius, 1898) in the coastal waters of

- Ondo state, Nigeria. Agric Sci Technol. 2021;13(4):425-31.
10. Udoh DI, Udo IU, Udoh EI. Microbiological analysis of the freshwater clam (*Galatea paradoxa*, Born 1778) caught from Cross River, Nigeria. Niger J Agric Food Environ. 2017;13(3):59-64.
 11. Rashed SM, Hasan NA, Alam M, Sadique A, Sultana M, Hoq MM et al. *Vibrio cholerae* O1 with reduced susceptibility to ciprofloxacin and azithromycin isolated from a rural coastal area of Bangladesh. Front Microbiol. 2017;8:252.
 12. Costa RA. *Escherichia coli* in sea food: A brief overview. Adv Biosci Biotechnol. 2013;4:450-4.
 13. Hariharan H, Amadi V. Shellfish as reservoirs of bacterial pathogens. J Coast Life Med. 2016;4:6-13.
 14. Orji JO, Nnachi AU, Ekwuatu CC, Akujobi CN, Iwuafor AA, Efunshile AM... Aghanya IN. Bacteriological quality of vended fresh shrimps harvested from Ndibe beach, Afikpo north local government area, Ebonyi state, Nigeria. Scholars Journal of Applied Medical Sciences (SJAMS). 2016;4(11C): 4058-4063. DOI: 10.21276/sjams.2016.4.11.40.
 15. Etang UE, Ikon GM, Udofia SM, Umo AN, Udo EE, Uyanga FZ, Ohagim PI. Microbiological analyses of kunu drinks locally produced and sold in Calabar, southern Nigeria. Journal of Advances in Microbiology. 2017;5(2):1-8. DOI: 10.9734/JAMB/2017/36169.
 16. Center for Food Safety, Food and Environmental Hygiene Department. Microbiological guidelines for food. Available at: https://www.cfs.gov.hk/english/food_leg/files/food_leg_Microbiological_Guidelines_for_Food_e.pdf. 2014; Accessed: Nov. 20th, 2022.
 17. Amin AN, Ahemd AM, Ahmed OM. Chemical and bacteriological risks of shrimp and clams (*Gandofly*) from Suez Gulf. Food Res. 2021;5(3):281-8. DOI: 10.26656/fr.2017.5(3).635.
 18. Talukder B, Chakma F, Sabuj AA-M, Haque ZF, Rahman T, Saha S. Identification, prevalence and antimicrobial susceptibility of major pathogenic bacteria in exportable shrimp (*Penaeus monodon*) of southern Bangladesh. Int J Fish Aquat Stud. 2019;7(1):96-9.
 19. Rosemary NO, Omokaro O, Wemedo SA, Ogbonna DN. Microbiological and nutritional quality of market and home smoke dried shrimp. Adv Microbiol. 2020; 20(5):35-42. DOI: 10.9734/JAMB/2020/v20i530243.
 20. Adebayo-Tayo AC, Odu NN, Michael MU, Okonko IO. Multi-drug resistant (MDR) organisms isolated from sea-foods in Uyo, south-southern Nigeria. Nat Sci. 2012;10:61-70.
 21. Gram L, Dalgaard P. Fish spoilage bacteria - Problems and solutions. Curr Opin Biotechnol. 2002;13(3):262-6. DOI: 10.1016/s0958-1669(02)00309-9
 22. Nwachukwu MI, Duru MKC, Odika PC, Udujih GO. Microbial evaluation of Njaba river and its effect on body weight of rats following consumption. J Chem Biol Phys Sci. 2013;3:2851-7.
 23. Moussa D, Bouba G, Daniel EM, Serge HZ, Moise N, Thomas N. Survival and growth of *Vibrio cholerae*, *Escherichia coli* and *Salmonella* spp in well water used for drinking purposes in Garova north (Cameroon). Int J Bacteriol. 2013;1(3):248-55.
 24. Center for Disease Control and Prevention. Relevant Pathogens From A-Z: *Pseudomonas* species; 2015. Available: <http://www.bodesciencecenter.com>; [accessed Dec 5 2017].
 25. Markey B, Leonard F, Archambault M, Cullinane A, Maguire D. Clinical veterinary microbiology. 2nd ed. New York: Oxford University Press; 2013.
 26. Khatri N, Tyagi S. Influence of natural and anthropogenic factors on surface and groundwater quality in rural and urban areas. Front Life Sci. 2015;8(1):23-39. DOI: 10.1080/21553769.2014.933716.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97614>