



Optimizing the Process Parameters for Carrot Cake Development Using Response Surface Methodology

**Chintan Prajapati ^{a*}, Keshav Kamaliya ^b
and Rashmin Dhingani ^a**

^a Department of Food Safety and Quality Assurance, College of Food Processing Technology and Bio-energy, Anand Agricultural University, Anand, Gujarat, India.

^b Polytechnic in Food Science and Home Economics, Anand Agricultural University, Anand, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ejns/2024/v16i91545>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/122398>

Original Research Article

Received: 14/07/2024

Accepted: 16/09/2024

Published: 21/09/2024

ABSTRACT

Carrots (*Daucus carota* L.) are rich in vitamins B1, B2, B6, β -carotene, and dietary fibers, having nutritional benefits such as, antioxidant, and anticancer properties. Carrot offers a valuable nutritional boost for enhancing bakery products. Incorporating carrot into cake preparation not only adds a unique flavor and moist texture but also boosts the nutritional value with additional vitamins and fiber. The research systematically examines the effects of varying process parameters such as carrot, sweetened condensed milk, and whey protein concentrate (WPC) on specific volume, springiness, and overall acceptability of carrot cake using Central Composite Design (CCD) of

*Corresponding author: Email: prajapatichintans5312@gmail.com;

Response Surface Methodology (RSM). The optimal formulation for carrot cake consisted of 70.8 g of carrot, 200 g of sweetened condensed milk, and 4.2 g of WPC. This combination resulted in a high-quality carrot cake with a specific volume of 2.58 cm³/g, a springiness measurement of 1.049 mm, and an overall acceptability rating of 8.3. This approach not only enhances product quality but also supports waste reduction by valorizing carrot, aligning with sustainable food processing practices.

Keywords: Carrot cake; response surface methodology; sweetened condensed milk; whey protein powder.

1. INTRODUCTION

The Indian bakery market crossed \$7.22 billion in 2018 globally and is projected to reach \$12 billion by 2024, with cakes expected to reach \$882 million. The industry includes traditional and modern products like breads, biscuits, and cakes. Increasing consumer demand for convenient, nutritious, and functional baked goods is responsible driving force for industrial growth [1].

The bakery industry can grow by developing fortified, nutritionally enhanced products. Bakery products, which are mainly made with wheat flour, can be enriched with health-promoting ingredients that contains omega-3 oils and dietary fibers. This may help prevent cardiovascular problems like atherosclerosis and irregular heartbeats, and may also help lower blood pressure. Adding dietary fibers may also be beneficial for individuals with diabetes as it can positively regulate blood glucose and cholesterol levels and promote gut health [2].

Carrots (*Daucus carota* L.) are the esteemed root vegetable of the *Apiaceae* family. Presently, Asia leads in carrot production at 61% [3]. In recent years, the consumption of carrots and their related products have increased consistently due to the recognition of antioxidant and anticancer activities of β -carotene in carrots, which is also a precursor of vitamin A [4]. Carrot also contains substantial nutrients such as proteins, lipids, sugars, vitamins, minerals, and dietary fibers [5]. These aforementioned nutritional benefits of carrots make it an excellent ingredient for enhancing bakery products, particularly cake which is the bakery product explored in this study. Carrots add vitamins (like vitamin A), fiber, and antioxidants to the cake, making it a slightly healthier option compared to a traditional wheat-based cake that might not offer these additional nutrients. In one research study, carrot pulp was used to prepare kulfi, that resulting in a safe, cost-effective, and consumer-friendly dessert that

is high in fiber and low in calories [6]. In a similar study, Turkish Delight was produced using carrot, orange, and carob pulp, leading to higher values of ash content, total phenolic substances, and antioxidant activity [7].

2. MATERIALS AND METHODS

2.1 Materials

The raw materials used in the production of carrot cake were procured from the local market of Anand, Gujarat. These include refined wheat flour (all-purpose flour), carrots (*Pusa yamdagni*), sweetened condensed milk, whey protein concentrate (WPC), ghee (shortening), sunflower oil, baking soda, baking powder, vinegar, essence, cake gel, skimmed milk powder (SMP), sugar, and PET container. The raw materials were of good quality and from the same reputable brand. They were purchased in one or as few instances as possible, stored in a cool, dry place, and used as required.

2.2 Methods

The methods for standardization of carrot cake are discussed in the following sections.

2.2.1 Processing of carrot

The raw carrots were cleaned, peeled, and grated. After that, the carrots were steam blanched at 90°C for 3 min. The blanched carrots were then chilled in a refrigerator at 5°C for 15min until the core temperature of the carrot slices reached 5°C. The chilled carrots were then grinded and packed in PE pouches, frozen and stored at -24°C [8].

2.2.2 Standardization of process parameters for carrot cake

The standardization of process parameters for carrot cake was carried out to establish

appropriate proportions of ingredients as shown in Table 1.

Table 1. Recipes for manufacturing of carrot cake

Ingredients	Amount
Refined wheat flour (g)	100
Ghee (g)	25
Sunflower oil (g)	25
Sweetened condensed milk(g)	50, 100, 150, 200, 250
WPC (g)	2, 4, 6, 8, 10
Carrot (g)	0, 25, 50, 75, 100
Water (g)	0, 25, 50, 75, 100
Baking soda (g)	1
Baking powder (g)	2
Vinegar (g)	6
Essence (g)	0.5
Cake gel (g)	3

2.2.3 Experimental design for carrot cake

A central composite design (CCD) was carried out to examine the effects of various responses influenced by three independent variables namely carrot (A), sweetened condensed milk (B), and whey protein concentrate (C) having 5 levels. The dependent variables were specific volume, springiness, and overall acceptability. Table 2 provides the variation levels of the independent and dependent variables. The levels were determined using coded values of -2, -1, 0, +1, and +2. The actual and coded values for each experimental combination were detailed in Table 3.

Table 2. Actual and coded values of three independent variables and dependent variables used in CCD

Independent variables	-Alpha	Low	Centre	High	+Alpha
	-2	-1	0	+1	+2
Carrot (g)	0	25	50	75	100
sweetened Condensed milk (g)	50	100	150	200	250
Whey protein concentrate (g)	2	4	6	8	10
Dependent variables	Observations				
Physical properties	Specific volume				
Textural characteristics	Springiness				
Sensory properties	Overall acceptability				

The carrot cake was prepared according to the procedure and the trials suggested by the software (Table 3). The optimization of independent variables was conducted using advanced statistical software, Design Expert 13.0.5.0. All 20 runs were executed and the results were statistically analyzed. Each of the 20 carrot cake samples were prepared with varying compositions as per the design and evaluated for physical, textural, and sensory properties. The data was analyzed using response surface methodology (RSM) and the optimization of independent variables was performed based on the recommended results by the software.

2.2.4 Process for manufacture of carrot cake

The carrot cake was made by first combining necessary ingredients, including sweetened condensed milk, whey protein concentrate (WPC), and cake gel. After that essence were added, then the mixture was blended with sunflower oil and ghee, and then water was incorporated. Refined wheat flour, baking powder, and baking soda were then mixed and added to the prepared mixture. Lastly, vinegar and carrot were incorporated into the batter, which was then poured into a cake mold and baked at 175°C for 50-55 minutes. After baking, the cake was cooled for 1-1.5 hours before being packed.

2.2.5 Analytical methods for proximate composition

The moisture, protein, fat, ash, crude fiber, and carbohydrates content of carrot cake was analyzed [9-11].

Table 3. Experimental design matrix for optimization of carrot cake using CCD of RSM

Run No.	Carrot (g)	Sweetened condensed milk (g)	WPC (g)
1	25	100	8
2	75	200	8
3	75	100	4
4	50	150	6
5	75	200	4
6	0	150	6
7	75	100	8
8	50	150	6
9	50	50	6
10	50	150	6
11	50	250	6
12	50	150	2
13	25	200	8
14	50	150	10
15	100	150	6
16	50	150	6
17	25	100	4
18	50	150	6
19	25	200	4
20	50	150	6

3. RESULTS AND DISCUSSION

3.1 Optimization of Process Parameters for Manufacturing of Carrot Cake Using RSM

This experiment aims to standardize the process parameters for manufacturing of carrot cake. The independent variables/ factors selected for optimization of the process parameters were carrot, sweetened condensed milk, and whey protein concentrate (WPC) and responses were specific volume, springiness, and overall acceptability is illustrated in Table 4. All the 20 runs were conducted according to the procedure shown in section 2.2.4 and evaluated for specific volume, springiness, and overall acceptability.

3.1.1 Influence of independent variables on properties of carrot cake

The responses obtained by conducting 20 runs, as presented in Tables 4, were individually analyzed using the quadratic regression model in the Design Expert software. ANOVA of the model was utilized to assess the significance of the model and the impact of individual factors on the responses. The fitted quadratic regression model for each response and their respective effects are discussed below.

3.1.2 Effect of process variables on specific volume of carrot cake

Specific volume is expressed as the volume per unit mass of a cake. The specific volume of carrot cakes varied within the range of 1.55 to 2.58 cm³/g as shown in Table 4. The specific volume was observed maximum (2.58 cm³/g) for the run 11 executed using 50 g carrot, 250 g sweetened condensed milk, and 6 g WPC while the minimum (1.55 cm³/g) for the run 9, with 50 g carrot, 50 g sweetened condensed milk, and 6 g WPC in Table 4.

As shown in Table 5, the quadratic model for specific volume achieved an F-value of 80.14 and a P-value of < 0.0001 indicating that the model was highly significant. The lack of Fit was not significant compared to the pure error, suggesting that the model was well fitted to the experiment. The multiple regression equation developed to predict the specific volume under the influence of various factors is presented below:

$$\text{Specific volume (cm}^3\text{/g)} = 2.45 + 0.1669^*A + 0.2356^*B - 0.0219^*C + 0.1313^*AB + 0.0087^*AC + 0.0087^*BC - 0.0699^*A^2 - 0.0886^*B^2 + 0.0014^*C^2$$

The values shown in Table 5 and above equation revealed that the linear parameters (A), and (B) shown a highly significant effect (P<0.01) on the




Fig. 1. Interaction of process variables on specific volume of carrot cake






Fig. 2. Interaction of process variables on springiness of carrot cake




Fig. 3. Interaction of process variables on overall acceptability of carrot cake

Table 4. Factors and responses of experimental carrot cake

Run No.	Factors			Responses		
	A	B	C	Specific volume (cm ³ /g)	Springiness (mm)	Overall acceptability (out of 9)
1	25	100	8	1.76	0.869	6.1
2	75	200	8	2.55	1.030	7.8
3	75	100	4	2.40	0.870	7.0
4	50	150	6	2.40	0.985	7.6
5	75	200	4	2.57	1.040	8.3
6	0	150	6	1.80	1.005	6.7
7	75	100	8	2.39	0.834	6.9
8	50	150	6	2.46	0.999	7.5
9	50	50	6	1.55	0.603	5.1
10	50	150	6	2.41	0.988	7.7
11	50	250	6	2.58	1.055	8.1
12	50	150	2	2.48	1.005	7.7
13	25	200	8	2.49	1.038	7.6
14	50	150	10	2.37	0.980	6.5
15	100	150	6	2.48	0.976	7.8
16	50	150	6	2.47	0.987	7.7
17	25	100	4	1.85	0.937	6.7
18	50	150	6	2.41	0.999	7.4
19	25	200	4	2.50	1.050	7.8
20	50	150	6	2.47	0.998	7.6

specific volume while, the effects of parameters (C) were not-significant ($P>0.05$). The quadratic term of the factor represents the impact of the ingredient at its highest level used during product standardization. The interaction effects of factors (AB) was highly significant ($P<0.01$) on the specific volume. The effect of the factor (A^2), and (B^2) was highly significant ($P<0.01$) while the (AC), (BC), and (C^2) parameters shown not-significant ($P>0.05$) effects at quadratic levels.

As indicated in Table 6, the adjusted R^2 and predicted R^2 for the specific volume were 0.97 and 0.91, respectively while the coefficient of determination was 0.98. A higher R^2 values approaching 1.00 indicated a better fit for the quadratic model. The adequate precision value (APV) for specific volume was 26.96, significantly surpassing the minimum desirable APV value of 4.00. This suggests that the model is reliable for predicting the response within the design space.

The three-dimensional response surfaces are plotted to illustrate the interactions among the variables on specific volume are shown in Fig. 1.

The effect of sweetened condensed milk and carrot on specific volume at constant 6 g WPC is depicted in Fig. 1. The results indicated that as the levels of carrot and sweetened condensed

milk increased, the specific volume also increased. The impact of sweetened condensed milk was more pronounced than that of carrot. The specific volume observed maximum (2.58 cm³/g) for the run 11 executed using 50 g carrot, and 250 g sweetened condensed milk at constant 6 g WPC while the minimum (1.55 cm³/g) for the run 9 executed with 50 g carrot, and 50 g sweetened condensed milk at constant 6 g WPC. Similar studies [2,12] observed that increasing the level of sweetened condensed milk increased the cake specific volume.

RSM plot (Fig. 1), showed the effect of WPC and carrot on specific volume at constant sweetened condensed milk (150 g). It was observed that as the level of carrot increased, the specific volume also increased, whereas increase in WPC level results in decreased specific volume. With sweetened condensed milk held constant at 150 g, the highest specific volume, 2.48 cm³/g, was observed in two experiments run number 12 with 50 g carrot and 2 g WPC, and in run number 15 with 100 g carrot and 6 g WPC. The minimum specific volume of 1.8 cm³/g was obtained in experiment run number 6, with a constant sweetened condensed milk level of 150 g, using 0 g carrot and 6 g WPC.

They observed that the increase the level of WPC, results in decrease cake's volume and specific volume [2].

Table 5. ANOVA table for effect of process variables on specific volume of carrot cake

Source	Sum of Squares	df	Mean Square	F-value	p-value	
Model	1.77	9	0.1966	80.14	< 0.0001	significant
A-Carrot	0.4456	1	0.4456	181.58	< 0.0001*	
B- Sweetened condensed milk	0.8883	1	0.8883	362.01	< 0.0001*	
C-WPC	0.0077	1	0.0077	3.12	0.1078	
AB	0.1378	1	0.1378	56.16	< 0.0001*	
AC	0.0006	1	0.0006	0.2496	0.6282	
BC	0.0006	1	0.0006	0.2496	0.6282	
A ²	0.1228	1	0.1228	50.04	< 0.0001*	
B ²	0.1975	1	0.1975	80.50	< 0.0001*	
C ²	0.0000	1	0.0000	0.0191	0.8930	
Residual	0.0245	10	0.0025			
Lack of Fit	0.0190	5	0.0038	3.43	0.1009	not significant

Significant at 1.0 percent level ($P < 0.01$), **significant at 5.0 percent level ($P < 0.05$)

Table 6. Fit statistics data of specific volume

Std. Dev.	0.05	R ²	0.98
Mean	2.32	Adjusted R ²	0.97
C.V. %	2.14	Predicted R ²	0.91
		APV	26.96

The effect of WPC and sweetened condensed milk on specific volume at constant carrot (50 g) is depicted in Fig. 1. It was observed that as the level of sweetened condensed milk increased, the specific volume also increased, whereas increase in WPC level results in decreased specific volume. With carrot held constant at 50 g, the highest specific volume, 2.58 cm³/g, was observed in experiment run number 11 with 250 g sweetened condensed milk and 6 g WPC. The minimum specific volume of 1.55 cm³/g was obtained in experiment run number 9, with a constant carrot level of 50 g, using 50 g sweetened condensed milk, and 6 g WPC.

3.1.3 Effect of process variables on springiness of carrot cake

The springiness of carrot cake varied within the range of 0.603 to 1.055 mm as shown in Table 4. The springiness was observed maximum of 1.055 mm for the run 11 executed using 50 g carrot, 250 g sweetened condensed milk, and 6 g WPC while the minimum of 0.603 mm for the run 9, with 50 g carrot, 50 g sweetened condensed milk, and 6 g WPC in Table 4. As shown in Table 7, the quadratic model for springiness achieved an F-value of 37.34 and a P-value of < 0.0001 indicating that the model was highly significant. The lack of Fit was not significant compared to the pure error, suggesting that the model was well fitted to the experiment. The multiple regression

equation developed to predict the springiness under the influence of various factors is presented below:

$$\text{Springiness (mm)} = 0.998 - 0.0111 * A + 0.097 * B - 0.011 * C + 0.0104 * AB + 0.0043 * AC + 0.0103 * BC - 0.001 * A^2 - 0.0414 * B^2 - 0.0005 * C^2$$

The values shown in Table 7 and above equation reveals that the linear parameters (B) shown a highly significant effect ($P < 0.01$) on the springiness while, the effects of parameters (A), and (C) were not-significant ($P > 0.05$). The quadratic term of the factor represents the impact of the ingredient at its highest level used during product standardization. The effect of the factor (B²) was highly significant ($P < 0.01$) while the (AB), (AC), (BC), (A²) and (C²) parameters shown not-significant ($P > 0.05$) effects at quadratic levels.

As indicated in Table 8, the adjusted R² and predicted R² for the specific volume were 0.94 and 0.79, respectively while the coefficient of determination (R²) was 0.97. A higher R² values approaching 1.00 indicated a better fit for the quadratic model. The adequate precision value (APV) for springiness was 24.17, significantly surpassing the minimum desirable APV value of 4.00. This suggests that the model is reliable for predicting the response within the design space.

The three-dimensional response surfaces are plotted to illustrate the interactions among the variables on springiness are shown in Fig. 2.

The effect of sweetened condensed milk and carrot on springiness at constant WPC (6 g) is depicted in Fig. 2. The results indicates that as the levels of sweetened condensed milk increased, the springiness also increased, whereas no significant effect of increase carrot level on springiness. The springiness was observed maximum (1.055 mm) for the run 11 executed using 50 g carrot, and 250 g sweetened condensed milk at constant 6 g WPC while the minimum of 0.603 mm for the run 9, with 50 g carrot, 50 g sweetened condensed milk at constant 6 g WPC. A similar study [12] observed that increasing the level of sweetened condensed milk increased the cake springiness.

RSM plot (Fig. 2), shows the effect of WPC and carrot on springiness at constant sweetened condensed milk (150 g). It was observed that as the levels of carrot and WPC increased, there was no significant effect on springiness. With sweetened condensed milk held constant at 150 g, the highest springiness of 1.023 mm was observed in experiment run number 20 with 50 g carrot and 6 g WPC. The minimum springiness of 0.976 mm was obtained in experiment run

number 15, with a constant sweetened condensed milk level of 150 g, using 100 g carrot and 6 g WPC.

The effect of WPC and sweetened condensed milk on springiness at constant carrot (50 g) is depicted in Fig. 2. It was observed that as the level of sweetened condensed milk increased, the springiness also increased, whereas no significant effect of increase WPC level on springiness. With carrot held constant at 50 g, the highest springiness of 1.055 mm was observed in experiment run number 11 with 250 g sweetened condensed milk and 6 g WPC. The minimum springiness of 0.603 mm was obtained in experiment run number 9, with a constant carrot level of 50 g, using 50 g sweetened condensed milk, and 6 g WPC.

3.1.4 Effect of process variables on overall acceptability of carrot cake

The overall acceptability of carrot cakes varied within the range of 5.1 to 8.3 (on 9-point hedonic scale base) as shown in Table 4. The overall acceptability was observed maximum (8.3) for the run 5 executed using 75 g carrot, 200 g sweetened condensed milk, and 4 g WPC while the minimum (5.1) for the run 9, with 50 g carrot, 50 g sweetened condensed milk, and 6 g WPC in Table 4.

Table 7. ANOVA table for effect of process variables on springiness of carrot cake

Source	Sum of Squares	df	Mean Square	F-value	p-value	
Model	0.2022	9	0.0225	37.34	< 0.0001*	significant
A-Carrot	0.0020	1	0.0020	3.27	0.1007	
B-Sweetened condensed milk	0.1504	1	0.1504	250.02	< 0.0001*	
C-WPC	0.0019	1	0.0019	3.23	0.1027	
AB	0.0009	1	0.0009	1.45	0.2570	
AC	0.0001	1	0.0001	0.2487	0.6288	
BC	0.0008	1	0.0008	1.40	0.2635	
A ²	0.0000	1	0.0000	0.0422	0.8414	
B ²	0.0431	1	0.0431	71.56	< 0.0001*	
C ²	6.401E-06	1	6.401E-06	0.0106	0.9199	
Residual	0.0060	10	0.0006			
Lack of Fit	0.0050	5	0.0010	4.96	0.0517	not significant

Significant at 1.0 percent level ($P < 0.01$), **significant at 5.0 percent level ($P < 0.05$)

Table 8. Fit statistics data of springiness

Std. Dev.	0.02	R ²	0.97
Mean	0.96	Adjusted R ²	0.94
C.V. %	2.55	Predicted R ²	0.79
		APV	24.17

Table 9. ANOVA table for effect of process variables on overall acceptability of carrot cake

Source	Sum of Squares	df	Mean Square	F-value	p-value	
Model	10.67	9	1.19	29.77	< 0.0001	significant
A-Carrot	1.0000	1	1.0000	25.10	0.0005*	
B- Sweetened condensed milk	7.29	1	7.29	182.98	< 0.0001*	
C-WPC	0.9025	1	0.9025	22.65	0.0008*	
AB	0.0200	1	0.0200	0.5020	0.4948	
AC	0.0050	1	0.0050	0.1255	0.7305	
BC	0.0000	1	0.0000	0.0000	1.0000	
A ²	0.1248	1	0.1248	3.13	0.1072	
B ²	1.36	1	1.36	34.25	0.0002*	
C ²	0.2930	1	0.2930	7.35	0.0219**	
Residual	0.3984	10	0.0398			
Lack of Fit	0.3301	5	0.0660	4.83	0.0544	not significant

Significant at 1.0 percent level ($P < 0.01$), **significant at 5.0 percent level ($P < 0.05$)

Table 10. Fit statistics data of overall acceptability

Std. Dev.	0.20	R ²	0.96
Mean	7.28	Adjusted R ²	0.93
C.V. %	2.74	Predicted R ²	0.76
		APV	20.96

As shown in Table 9, the quadratic model for overall acceptability achieved an F-value of 29.77 and a P-value of < 0.0001 indicating that the model was highly significant. The lack of Fit was not significant compared to the pure error, suggesting that the model was well fitted to the experiment. The multiple regression equation developed to predict the overall acceptability under the influence of various factors is presented below:

$$\text{Overall acceptability} = 7.61 + 0.25^* A + 0.675^* B - 0.2375^* C - 0.05^* AB + 0.025^* AC + 0^* BC - 0.0705^* A^2 - 0.233^* B^2 - 0.108^* C^2$$

The values shown in Table 9 and above equation revealed that the linear parameters (A), (B), and (C) showed a highly significant effect ($P < 0.01$) on the overall acceptability.

The quadratic term of the factor represents the impact of the ingredient at its highest level used during product standardization. The effect of the factor (B²) was highly significant ($P < 0.01$). The interaction effects of factors (C²) was significantly ($P < 0.05$) on the overall acceptability as shown in Table 9 while the (AB), (AC), (BC), and (A²) parameters shown not-significant ($P > 0.05$) effects at quadratic levels.

As indicated in Table 10, the adjusted R² and predicted R² for the overall acceptability were

0.93 and 0.76, respectively while the coefficient of determination (R²) was 0.96. A higher R² values approaching 1.00 indicated a better fit for the quadratic model. The adequate precision value (APV) for overall acceptability was 20.96, significantly surpassing the minimum desirable APV value of 4.00. This suggests that the model is reliable for predicting the response within the design space.

The three-dimensional response surfaces are plotted to illustrate the interactions among the variables on overall acceptability are shown in Fig. 3.

The effect of sweetened condensed milk and carrot on overall acceptability at constant WPC (6 g) is depicted in Fig. 3. The results indicated that as the levels of carrot and sweetened condensed milk increased, the overall acceptability also increased. It was observed that, the impact of sweetened condensed milk was more pronounced than that of carrot. The overall acceptability was observed maximum of 8.1 for the run 11 executed using 50 g carrot, and 250 g sweetened condensed milk at constant 6 g WPC while the minimum (5.1) for the run 9, with 50 g carrot, and 50 g sweetened condensed milk at constant 6 g WPC.

RSM plot (Fig. 3), showed the effect of WPC and carrot on overall acceptability at constant sweetened condensed milk (150 g). It was observed that as the level of carrot increased, the overall acceptability also increased, whereas increase in WPC level results in decreased overall acceptability. With sweetened condensed milk held constant at 150 g, the highest overall acceptability of 7.8 was observed in experiment run number 15 with 100 g carrot and 6 g WPC. The minimum overall acceptability of 6.5 was obtained in experiment run number 14, with a constant sweetened condensed milk level of 150 g, using 50 g carrot and 10 g WPC.

The effect of WPC and sweetened condensed milk on overall acceptability at constant carrot (50 g) is depicted in Fig. 3. It was observed that as the level of sweetened condensed milk increased, the overall acceptability also increased, whereas increase in WPC level results in decreased overall acceptability. In experiment run number 11 where in 250 g sweetened condensed milk and 6 g WPC were added with carrot held constant at 50 g, the highest overall acceptability of 8.1 was observed.

The minimum overall acceptability of 5.1 was obtained in experiment run number 9, with a constant carrot level of 50 g, using 50 g sweetened condensed milk, and 6 g WPC. Similar studies [13-15] observed that increasing the level of carrot and sweetened condensed milk increased the cake overall acceptability.

3.1.5 Optimization of process parameters for carrot cake using quadratic regression model of response surface methodology

The optimization of process parameters was done to find the optimal combinations of variables for making carrot cake viz. carrot, sweetened condensed milk, and WPC that would result in the product having the most acceptable specific volume, springiness, and overall acceptability.

The data received for the responses were examined in Design-Expert Software. The optimization criteria/targets were as follows, as shown in Table 11.

Table 11. Selected criteria/ targets for optimization of process parameters for carrot cake

Name	Criteria/ targets	Lower Limit	Upper Limit
Independent Variables			
Carrot (g)	is in range	25	75
Sweetened condensed milk (g)	is in range	100	200
WPC (g)	is in range	4	8
Dependent Variables			
Specific volume (cm ³ /g)	maximize	1.55	2.58
Springiness (mm)	maximize	0.603	1.055
Overall Acceptability (out of 9)	maximize	5.1	8.3

Table 12. Optimized solution for carrot cake as suggested by quadratic regression model of RSM

Optimized Factors			Optimized Responses			Desirability of the model
Carrot (g)	Sweetened condensed milk (g)	WPC (g)	Specific volume (cm ³ /g)	Springiness (mm)	Overall acceptability (out of 9)	0.99
70.8	200	4.2	2.58	1.049	8.3	

Table 13. Validation of the optimized RSM model for carrot cake

Analysis	Predicted value	95% PI low	Actual value	95% PI high
Specific volume (cm ³ /g)	2.58	2.48	2.59 ± 0.01	2.68
Springiness (mm)	1.049	1.00	1.046 ± 0.01	1.10
Overall acceptability (out of 9)	8.28	7.90	8.10 ± 0.1	8.60

Values expressed as mean ± sd (n=3)

Using the parameters and their limits, the RSM recommended an optimized solution, which is shown in Table 12. That includes 70.8 g carrot, 200 g sweetened condensed milk, and 4.2 g WPC and responses were 2.58 cm³/g specific volume, 1.049 mm springiness, and 8.3 overall acceptability as the model was 99% desirable.

Table 13 shows the validation of the carrot cake of the actual values for specific volume (2.59 ± 0.01 cm³/g), springiness (1.046 ± 0.01 mm), and overall acceptability (8.10 ± 0.1) closely align with the predicted values, remaining well within their respective 95% prediction intervals.

3.2 Proximate Composition of Optimized Carrot Cake

The developed carrot cake has 27.49, 7.01, 15.6, 0.01, 1.57 and 49.89% moisture, protein, fat, ash, crude fiber, and carbohydrates (by difference), respectively.

4. CONCLUSION

The present research had conducted in order to highlight the benefits of carrot into the bakery products in which the effect of process parameters such as carrot, sweetened condensed milk, and whey protein concentrate (WPC) on the quality attributes of cake, including specific volume, springiness, and overall acceptability was systematically assessed using CCD of RSM. Interaction effect of process variables was evaluated at five levels with 20 run experimental design. The quadratic model was suggested by software and showed a well fit for the specific volume, springiness, and overall acceptability of carrot cake, with R² values of 0.98, 0.97, and 0.96, respectively. The quadratic model for specific volume had an F-value of 80.14 and a P-value of < 0.0001, indicating highly significance and the lack of fit was not significant, confirming the model fitted well. Linear parameters (A) and (B) had a significant impact (P<0.01) on specific volume, while parameter (C) was not significant (P>0.05). Furthermore, the interaction of factors (AB) was highly significant (P<0.01) and quadratic effects (A²) and (B²) were significant (P<0.01). However, (AC), (BC), and (C²) were not significant (P>0.05). The quadratic model for springiness had an F-value (37.34) and P < 0.0001, showing higher significance level and the lack of fit was not significant, confirming a good model fit. Linear parameter (B) had a significant effect (P<0.01) on springiness, while parameters (A)

and (C) were not significant (P>0.05). The quadratic effect of (B²) was highly significant (P<0.01), whereas interactions (AB), (AC), (BC), and quadratic terms (A²) and (C²) were not significant (P>0.05). A quadratic model showed higher significance level for overall acceptability having F-value (29.77) and P < 0.0001, indicating well-fitted model without significant lack of fit. Linear parameters (A, B, C) significantly affected the acceptability (P < 0.01), with the quadratic term (B²) showing highly significant effect (P < 0.01). Interaction effects (C²) were significant (P < 0.05), while (AB), (AC), (BC), and (A²) were not significant (P > 0.05) at quadratic levels. Based on response surface methodology (RSM), an optimized solution for carrot cakes was recommended in a blend of 70.8 g carrot, 200 g sweetened condensed milk, and 4.2 g WPC. The corresponding responses were specific volume (2.58 cm³/g), springiness (1.049 mm), and overall acceptability (8.3 (out of 9)). The model achieved 99% desirability, indicating highly favorable results within the predefined parameter limits. The carrot added significant nutritional benefits, including additional vitamins and dietary fiber, thereby enhancing the health profile of the final product. This approach might support healthier bakery options with contributing to more environmentally friendly food processing practices.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mehdi MM, Rakshit S, Sarma TR, Joshi M, Nille N. Entrepreneurship and innovation in the bakery industry: A case study of ganesh bakery, FIIB Business Review. 2022;11(1):30-35.
2. Zhou W, Hui YH, De Leyn I, Pagani MA, Rosell CM, Selman JD, Therdthai N. Bakery Products Science and Technology. John Wiley & Sons; 2014.

3. Krivokapić S, Pejatović T, Perović S. Chemical characterization, nutritional benefits and some processed products from carrot (*Daucus carota* L.). *Poljoprivreda i Sumarstvo*. 2020;66(2):191-216.
4. Reddy MNK, Kumar MS, Reddy GB, Reddy NA, Rao VK. Quality evaluation of turkey meat sausages incorporated with ground carrot. *The Pharma Innovation Journal*. 2018;7:773-777.
5. Bao B, Chang KC. Carrot pulp chemical composition, color, and water-holding capacity as affected by blanching. *Journal of Food Science*. 1994;59(6):1159-1161.
6. Akhter MJ, Kabir MI, Sohany M, Islam MH, Khatun AA, Hosen A, Kabir MF. Effect of carrot pulp on the physicochemical, microbiological and sensory attributes of kulfi. *Food Research*. 2024;8(4):1-9.
7. Hanoğlu A, Karaoğlu MM, Bedir Y. The effect of carob, orange and carrot pulps on physical, chemical and microbiological properties of Turkish delight. *International Journal of Gastronomy and Food Science*. 2023;32:100-709.
8. Kidmose U, Martens HJ. Changes in texture, microstructure and nutritional quality of carrot slices during blanching and freezing. *Journal of the Science of Food and Agriculture*. 1999;79(12):1747-1753.
9. AOAC International, Official methods of analysis of AOAC International. 2 vols 22nd ed. Rockville, MD: Association of Official Analytical Chemists; 2023.
10. Ranganna S, Handbook of analysis and quality control for fruit and vegetable products, The McGraw-Hill Education (India) Pvt. Ltd., New Delhi; 2019.
11. ICMR & NIN, Nutrient requirements for Indians: Carbohydrates. Department of health research, ministry of health & family welfare; Government of India; 2020.
12. Khetarpaul N, Grewal RB, Jood S. *Bakery Science and Cereal Technology*. Daya Books; 2016.
13. Amany EM, Youssef MM, Abou-Gharbia HA, Hamida MM. Utilization of carrot pomace in formulating functional biscuits and cakes. *Alex. Journal of Food Science and Technology*. 2010;7(2):25-32.
14. Austin PP. Development and quality evaluation of orange peel and carrot pomace powder incorporated cake [Doctoral dissertation, Sam Higginbottom University of Agriculture, Technology and Sciences, Allahabad, (UP), India]. Retrieved from; 2019. Available:<http://krishikosh.egranth.ac.in/handle/1/5810137218>
15. Mohtarami F. Effect of carrot pomace powder and dushab (traditional grape juice concentrate) on the physical and sensory properties of cakes: A combined mixtures design approach. *Current Nutrition & Food Science*. 2019;15(6):572-82

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/122398>